Floaters and Flashes






Overview

We have developed this leaflet to help you understand more about floaters and flashes.

Floaters look like small, dark spots or strands that appear to float in front of your eyes. Floaters are very common and are normally harmless. They are more common if you are short sighted or as you get older.

Some people notice they see **flashes** of light. These can be due to movement of the gel inside the eye.

Very occasionally, flashes or an increase in floaters can be a sign of a retinal detachment, which needs treating as soon as possible. This is more common as you get older, or in people who are short sighted or have had eye surgery. If you have any concerns about the health of your eyes please visit your local optometrist. Optometrists are the eye-health specialists on the high street. An eye examination is a vital health check and should be part of everyone's normal health care.



If you get any of the following symptoms and you cannot contact your optometrist, you should seek urgent attention ideally from an eye casualty department at the hospital. It is important that you seek advice promptly if you have:

- a sudden increase in floaters, particularly if you also notice flashing lights
- · a new, large, floater
- a change in floaters or flashing lights after you have had a direct blow to your eye
- a shadow spreading across the vision of one of your eyes.

What are floaters?

Floaters appear as black spots or something that looks like a hair or small pieces of a cobweb. These can be semi-transparent or dark and appear to float in front of your vision. If you have had these for years, your eye and your brain learn to ignore them. Sometimes the number of floaters increases as you get older. Occasionally an increase in floaters can be a sign of problems inside the eye.

Because they 'float' in the jelly of your eye, you will find that if you move your eye to try to look at a floater it will move away in the direction you move your eye. You might only see the floater if you are staring at a light coloured surface or at the sky during the day.

Some people find that floaters can be a nuisance, but most people become used to them. They rarely cause problems with your vision.

"Floaters appear as black spots or something that looks like a hair or small pieces of a cobweb"

Why do floaters occur?

Some people are born with floaters. Other floaters occur as you get older when the gel in the eye, the vitreous humour, naturally shrinks. The gel separates into a watery fluid and wavy collagen fibrils. The fibrils are seen as line-shaped floaters. Sometimes the gel shrinks enough to collapse away from the light sensitive lining at the back of your eye, which is called the retina. Once the gel has collapsed, some people see a large ring-shaped floater.

The collapse of the vitreous gel can pull on your retina. If this happens you would see this as flashes of light – see 'flashes', opposite.

Floaters can also be caused by some eye diseases that cause inflammation. This is not very common.

Who is at risk of floaters?

Floaters are more common in people who are short sighted. They may increase if you have had an eye operation such as cataract surgery, or laser treatment after cataract surgery.

What might happen if I have floaters?

Most of the time floaters are harmless. Sometimes they may be annoying, but treatment is not advised.

Occasionally a sudden increase in floaters – either one or more large ones or a shower of tiny ones – may be a sign of more serious eye

"A sudden increase in floaters – either one or more large ones or a shower of tiny ones – may be a sign of a more serious eye disease such as a retinal detachment"

disease such as a retinal detachment. This is when your retina pulls away from the back of your eye: it may lead to a sudden increase in floaters and possibly a blank spot or shadow in your vision which does not go away. This needs immediate attention. If you notice these symptoms you should contact your optometrist straight away. If you can not do this you should seek urgent attention from an eye casualty department at the hospital. If there is no eye casualty department nearby, you can go to your usual hospital casualty department, but it is best to go to an eye casualty department if you can. An ophthalmologist, a specialist eye doctor, will need to use eye drops and a special light to look inside your eyes to check if your retina is damaged.

What are flashes?

Some people may see flashes of light in front of one of their eyes, like small sparkles, lightning or fireworks. These tend to be in the extreme corners of your vision, come and go, and don't obscure any part of your vision. The flashes don't last for a defined length of time, and you may notice them more if you go from a light to dark environment. They may continue for several months. These are different to the shimmering or zig-zag lines that may be part of a migraine. Migraine shimmers are a flickering of light, often on only one side of your vision, which then expands to the outside of your vision with a sort of jagged pattern. This will often obscure at least

part of your vision (the left or right side). The shimmers usually go away after 10-20 minutes and may be followed by a headache, although some people may get migraine shimmers even if they do not have a headache afterwards.

Flashes occur when there is a pull on your retina. This might happen as the vitreous gel inside your eye becomes more liquid and collapses. You may experience flashes occasionally, on and off over weeks or months. Flashes can also occur if you are hit in the eye.

Flashes related to a collapse of the gel inside the eye are more likely to happen as you get older.

What might happen if I have flashes?

Sometimes flashes just indicate a tug on the retina and nothing more. However constant flashes may be a sign of a retinal detachment.

A retinal tear or retinal detachment may lead to a sudden increase in floaters as well as flashes. You might notice a shadow at the edge of your vision too. This needs immediate attention. If you notice these symptoms you should contact your optometrist straight away. If you can not do this you should seek urgent attention from an eye casualty department at the hospital. If there is no eye casualty department nearby you can go to your usual hospital casualty department, but it is best to go to an eye casualty department if you can. An ophthalmologist, a specialist eye doctor, will need to use eye drops and a special light

"Flashes occur when there is a pull on your retina"

"Constant flashes may be a sign of a retinal detachment" to look inside your eyes to check if your retina is damaged.

Who is at risk of a retinal detachment?

Some people are more at risk of a retinal detachment. These are people who:

- have had eye surgery, such as a cataract operation or laser surgery after a cataract operation
- are moderately short sighted (over -3.00D)
- · have had a previous eye injury
- have a family history of retinal detachment
- have had a previous retinal detachment in that eye or the other eye
- are over the age of 50
- have certain retinal diseases such as lattice or other retinal degeneration
- have certain systemic diseases such as Marfan syndrome.

For more information please talk to your local optometrist.

This leaflet is produced by the College of Optometrists. We are the professional, scientific and examining body for optometry in the UK. People who are our members agree to meet the highest clinical and ethical standards. Look for the letters MCOptom or FCOptom to see if your optometrist is a member.

Please visit www.lookafteryoureyes.org for more information.

Your local optometrist:

If you would like this leaflet in large print, please email patients@college-optometrists.org

The College of Optometrists 42 Craven Street London, WC2N 5NG Tel: 020 7839 6000

Email: optometry@college-optometrists.org www.college-optometrists.org www.lookafteryoureyes.org



© College of Optometrists 2011